

This Page Is Inserted by IFW Operations
and is not a part of the Official Record

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images may include (but are not limited to):

- BLACK BORDERS
- TEXT CUT OFF AT TOP, BOTTOM OR SIDES
- FADED TEXT
- ILLEGIBLE TEXT
- SKEWED/SLANTED IMAGES
- COLORED PHOTOS
- BLACK OR VERY BLACK AND WHITE DARK PHOTOS ✓
- GRAY SCALE DOCUMENTS

IMAGES ARE BEST AVAILABLE COPY.

**As rescanning documents *will not* correct images,
please do not report the images to the
Image Problem Mailbox.**




Figure 1. Typical workflow for current methodologies used to calculate LV functional parameters.




Figure 2. New workflow for accurate non-invasive measurement of cardiac function using tomographic images





Figure 3. Representative Short Axis, Vertical Long Axis, Horizontal Long Axis, and Left Ventricle Inflow/Outflow Tract views from a cardiac CT Exam




Figure 4. Representation of delineation of the LV from surrounding anatomy via thresholding, edge detection, and region growing algorithms. This is done at both end diastole and end systole to segment the contrast from the ventricular walls and ventricular walls from the contrast.





Figure 5. An optimal combination of advanced algorithms such as thresholding, morphological and connectivity tools, edge detection, and region growing are used to segment the contrast within the ventricle from the myocardium.





Figure 6. Examples of 3D models of the LV at both end systole and end diastole. By measuring the volume of these models, EDV, ESV, SV, EF, and CO are calculated.



Figure 7. Volume rendering of the LV contrast at both end diastole and end systole.




FIG. 8




FIG. 9




FIG. 10